

PLANT PHYSIOLOGY

Fifth Edition



Lincoln Taiz

Professor Emeritus

University of California, Santa Cruz

Eduardo Zeiger

Professor Emeritus

University of California, Los Angeles



**Sinuer Associates Inc., Publishers
Sunderland, Massachusetts U.S.A.**

Contents

CHAPTER 1 Plant Cells 1

Plant Life: Unifying Principles 2

Overview of Plant Structure 2

- Plant cells are surrounded by rigid cell walls 2
- New cells are produced by dividing tissues called meristems 2
- Three major tissue systems make up the plant body 4

Plant Cell Organelles 4

- Biological membranes are phospholipid bilayers that contain proteins 4

The Endomembrane System 8

- The nucleus contains the majority of the genetic material 8
- Gene expression involves both transcription and translation 10
- The endoplasmic reticulum is a network of internal membranes 10
- Secretion of proteins from cells begins with the rough ER (RER) 13
- Glycoproteins and polysaccharides destined for secretion are processed in the Golgi apparatus 14
- The plasma membrane has specialized regions involved in membrane recycling 16
- Vacuoles have diverse functions in plant cells 16

Independently Dividing Organelles Derived from the Endomembrane System 17

- Oil bodies are lipid-storing organelles 17
- Microbodies play specialized metabolic roles in leaves and seeds 17

Independently Dividing, Semiautonomous Organelles 18

- Proplastids mature into specialized plastids in different plant tissues 21
- Chloroplast and mitochondrial division are independent of nuclear division 21

The Plant Cytoskeleton 22

- The plant cytoskeleton consists of microtubules and microfilaments 22
- Microtubules and microfilaments can assemble and disassemble 23
- Cortical microtubules can move around the cell by "treadmilling" 24
- Cytoskeletal motor proteins mediate cytoplasmic streaming and organelle traffic 24

Cell Cycle Regulation 25

- Each phase of the cell cycle has a specific set of biochemical and cellular activities 26
- The cell cycle is regulated by cyclins and cyclin-dependent kinases 26
- Mitosis and cytokinesis involve both microtubules and the endomembrane system 27

Plasmodesmata 29

- Primary and secondary plasmodesmata help to maintain tissue developmental gradients 29

SUMMARY 31

CHAPTER 2 Genome Organization and Gene Expression 35

Nuclear Genome Organization 35

The nuclear genome is packaged into chromatin 36

Centromeres, telomeres, and nucleolar organizers contain repetitive sequences 36

Transposons are mobile sequences within the genome 37

Polyploids contain multiple copies of the entire genome 38

Phenotypic and physiological responses to polyploidy are unpredictable 41

Plant Cytoplasmic Genomes: Mitochondria and Chloroplasts 42

The endosymbiotic theory describes the origin of cytoplasmic genomes 42

Organelle genomes consist mostly of linear chromosomes 43

Organelle genetics do not obey Mendelian laws 44

Transcriptional Regulation of Nuclear Gene Expression 45

RNA polymerase II binds to the promoter region of most protein-coding genes 45

Epigenetic modifications help determine gene activity 48

Posttranscriptional Regulation of Nuclear Gene Expression 50

RNA stability can be influenced by *cis*-elements 50

Noncoding RNAs regulate mRNA activity via the RNA interference (RNAi) pathway 50

Posttranslational regulation determines the life span of proteins 54

Tools for Studying Gene Function 55

Mutant analysis can help to elucidate gene function 55

Molecular techniques can measure the activity of genes 55

Gene fusions can introduce reporter genes 56

Genetic Modification of Crop Plants 59

Transgenes can confer resistance to herbicides or plant pests 59

Genetically modified organisms are controversial 60

SUMMARY 61

UNIT I Transport and Translocation of Water and Solutes 65

CHAPTER 3 Water and Plant Cells 67

Water in Plant Life 67

The Structure and Properties of Water 68

Water is a polar molecule that forms hydrogen bonds 68

Water is an excellent solvent 69

Water has distinctive thermal properties relative to its size 69

Water molecules are highly cohesive 69

Water has a high tensile strength 70

Diffusion and Osmosis 71

Diffusion is the net movement of molecules by random thermal agitation 71

Diffusion is most effective over short distances 72

Osmosis describes the net movement of water across a selectively permeable barrier 73

Water Potential 73

The chemical potential of water represents the free-energy status of water 74

Three major factors contribute to cell water potential 74

Water potentials can be measured 75

Water Potential of Plant Cells 75

Water enters the cell along a water potential gradient 75

Water can also leave the cell in response to a water potential gradient 77

Water potential and its components vary with growth conditions and location within the plant 77

Cell Wall and Membrane Properties 78

XVIII TABLE OF CONTENTS

Small changes in plant cell volume cause large changes in turgor pressure 78

The rate at which cells gain or lose water is influenced by cell membrane hydraulic conductivity 79

Aquaporins facilitate the movement of water across cell membranes 79

Plant Water Status 80

Physiological processes are affected by plant water status 80

Solute accumulation helps cells maintain turgor and volume 80

SUMMARY 81

CHAPTER 4 Water Balance of Plants 85

Water in the Soil 85

A negative hydrostatic pressure in soil water lowers soil water potential 86

Water moves through the soil by bulk flow 87

Water Absorption by Roots 87

Water moves in the root via the apoplast, symplast, and transmembrane pathways 88

Solute accumulation in the xylem can generate “root pressure” 89

Water Transport through the Xylem 90

The xylem consists of two types of tracheary elements 90

Water moves through the xylem by pressure-driven bulk flow 92

Water movement through the xylem requires a smaller pressure gradient than movement through living cells 93

What pressure difference is needed to lift water 100 meters to a treetop? 93

The cohesion–tension theory explains water transport in the xylem 93

Xylem transport of water in trees faces physical challenges 94

Plants minimize the consequences of xylem cavitation 96

Water Movement from the Leaf to the Atmosphere 96

Leaves have a large hydraulic resistance 96

The driving force for transpiration is the difference in water vapor concentration 96

Water loss is also regulated by the pathway resistances 98

Stomatal control couples leaf transpiration to leaf photosynthesis 98

The cell walls of guard cells have specialized features 99

An increase in guard cell turgor pressure opens the stomata 101

The transpiration ratio measures the relationship between water loss and carbon gain 101

Overview: The Soil–Plant–Atmosphere Continuum 102

SUMMARY 102

CHAPTER 5 Mineral Nutrition 107

Essential Nutrients, Deficiencies, and Plant Disorders 108

Special techniques are used in nutritional studies 110

Nutrient solutions can sustain rapid plant growth 110

Mineral deficiencies disrupt plant metabolism and function 113

Analysis of plant tissues reveals mineral deficiencies 117

Treating Nutritional Deficiencies 117

Crop yields can be improved by addition of fertilizers 118

Some mineral nutrients can be absorbed by leaves 118

Soil, Roots, and Microbes 119

Negatively charged soil particles affect the adsorption of mineral nutrients 119

Soil pH affects nutrient availability, soil microbes, and root growth 120

Excess mineral ions in the soil limit plant growth 120

Plants develop extensive root systems 121

Root systems differ in form but are based on common structures 121

Different areas of the root absorb different mineral ions 123
 Nutrient availability influences root growth 124
 Mycorrhizal fungi facilitate nutrient uptake by roots 125

Nutrients move from mycorrhizal fungi to root cells 126

SUMMARY 126

CHAPTER 6 Solute Transport 131

Passive and Active Transport 132

Transport of Ions across Membrane Barriers 133

Different diffusion rates for cations and anions produce diffusion potentials 134
 How does membrane potential relate to ion distribution? 134
 The Nernst equation distinguishes between active and passive transport 136
 Proton transport is a major determinant of the membrane potential 137

Membrane Transport Processes 137

Channels enhance diffusion across membranes 139
 Carriers bind and transport specific substances 140
 Primary active transport requires energy 140
 Secondary active transport uses stored energy 142
 Kinetic analyses can elucidate transport mechanisms 143

Membrane Transport Proteins 144

The genes for many transporters have been identified 144
 Transporters exist for diverse nitrogen-containing compounds 146
 Cation transporters are diverse 147
 Anion transporters have been identified 148
 Metal transporters transport essential micronutrients 149
 Aquaporins have diverse functions 149
 Plasma membrane H⁺-ATPases are highly regulated P-type ATPases 150
 The tonoplast H⁺-ATPase drives solute accumulation in vacuoles 151
 H⁺-pyrophosphatases also pump protons at the tonoplast 153

Ion Transport in Roots 153

Solutes move through both apoplast and symplast 153
 Ions cross both symplast and apoplast 153
 Xylem parenchyma cells participate in xylem loading 154

SUMMARY 156

UNIT II Biochemistry and Metabolism 161

CHAPTER 7 Photosynthesis: The Light Reactions 163

Photosynthesis in Higher Plants 164

General Concepts 164

Light has characteristics of both a particle and a wave 164
 When molecules absorb or emit light, they change their electronic state 165
 Photosynthetic pigments absorb the light that powers photosynthesis 166

Key Experiments in Understanding Photosynthesis 167

Action spectra relate light absorption to photosynthetic activity 168

Photosynthesis takes place in complexes containing light-harvesting antennas and photochemical reaction centers 169
 The chemical reaction of photosynthesis is driven by light 170
 Light drives the reduction of NADP and the formation of ATP 171
 Oxygen-evolving organisms have two photosystems that operate in series 171

Organization of the Photosynthetic Apparatus 172

The chloroplast is the site of photosynthesis 172

XX TABLE OF CONTENTS

- Thylakoids contain integral membrane proteins 173
- Photosystems I and II are spatially separated in the thylakoid membrane 174
- Anoxygenic photosynthetic bacteria have a single reaction center 174

Organization of Light-Absorbing

Antenna Systems 176

- Antenna systems contain chlorophyll and are membrane associated 176
- The antenna funnels energy to the reaction center 176
- Many antenna pigment–protein complexes have a common structural motif 176

Mechanisms of Electron Transport 178

- Electrons from chlorophyll travel through the carriers organized in the “Z scheme” 178
- Energy is captured when an excited chlorophyll reduces an electron acceptor molecule 179
- The reaction center chlorophylls of the two photosystems absorb at different wavelengths 180
- The photosystem II reaction center is a multisubunit pigment–protein complex 181
- Water is oxidized to oxygen by photosystem II 181
- Pheophytin and two quinones accept electrons from photosystem II 183
- Electron flow through the cytochrome b_6f complex also transports protons 183
- Plastoquinone and plastocyanin carry electrons between photosystems II and I 184

- The photosystem I reaction center reduces NADP^+ 185
- Cyclic electron flow generates ATP but no NADPH 185
- Some herbicides block photosynthetic electron flow 186

Proton Transport and ATP Synthesis in the Chloroplast 187

Repair and Regulation of the Photosynthetic Machinery 189

- Carotenoids serve as photoprotective agents 190
- Some xanthophylls also participate in energy dissipation 190
- The photosystem II reaction center is easily damaged 191
- Photosystem I is protected from active oxygen species 191
- Thylakoid stacking permits energy partitioning between the photosystems 191

Genetics, Assembly, and Evolution of Photosynthetic Systems 192

- Chloroplast genes exhibit non-Mendelian patterns of inheritance 192
- Most chloroplast proteins are imported from the cytoplasm 192
- The biosynthesis and breakdown of chlorophyll are complex pathways 192
- Complex photosynthetic organisms have evolved from simpler forms 193

SUMMARY 194

CHAPTER 8 Photosynthesis: The Carbon Reactions 199

The Calvin–Benson Cycle 200

- The Calvin–Benson cycle has three stages: carboxylation, reduction, and regeneration 200
- The carboxylation of ribulose 1,5-bisphosphate fixes CO_2 for the synthesis of triose phosphates 201
- Ribulose 1,5-bisphosphate is regenerated for the continuous assimilation of CO_2 201
- An induction period precedes the steady state of photosynthetic CO_2 assimilation 204

Regulation of the Calvin–Benson Cycle 205

- The activity of rubisco increases in the light 206
- Light regulates the Calvin–Benson cycle via the ferredoxin–thioredoxin system 207

- Light-dependent ion movements modulate enzymes of the Calvin–Benson cycle 208
- Light controls the assembly of chloroplast enzymes into supramolecular complexes 208

The C_2 Oxidative Photosynthetic Carbon Cycle 208

- The carboxylation and the oxygenation of ribulose 1,5-bisphosphate are competing reactions 210
- Photorespiration depends on the photosynthetic electron transport system 213
- Photorespiration protects the photosynthetic apparatus under stress conditions 214
- Photorespiration may be engineered to increase the production of biomass 214

Inorganic Carbon–Concentrating Mechanisms 216**Inorganic Carbon–Concentrating Mechanisms: The C₄ Carbon Cycle 216**

Malate and aspartate are carboxylation products of the C₄ cycle 217

Two different types of cells participate in the C₄ cycle 218

The C₄ cycle concentrates CO₂ in the chloroplasts of bundle sheath cells 220

The C₄ cycle also concentrates CO₂ in single cells 221

Light regulates the activity of key C₄ enzymes 221

In hot, dry climates, the C₄ cycle reduces photorespiration and water loss 221

Inorganic Carbon–Concentrating Mechanisms: Crassulacean Acid Metabolism (CAM) 221

CAM is a versatile mechanism sensitive to environmental stimuli 223

Accumulation and Partitioning of Photosynthates—Starch and Sucrose 224**Formation and Mobilization of Chloroplast Starch 225**

Starch is synthesized in the chloroplast during the day 225

Starch degradation at night requires the phosphorylation of amylopectin 228

The export of maltose prevails in the nocturnal breakdown of transitory starch 230

Sucrose Biosynthesis and Signaling 231

Triose phosphates supply the cytosolic pool of three important hexose phosphates in the light 231

Fructose 2,6-bisphosphate regulates the hexose phosphate pool in the light 235

The cytosolic interconversion of hexose phosphates governs the allocation of assimilated carbon 235

Sucrose is continuously synthesized in the cytosol 235

SUMMARY 237**CHAPTER 9 Photosynthesis: Physiological and Ecological Considerations 243****Photosynthesis Is the Primary Function of Leaves 244**

Leaf anatomy maximizes light absorption 245

Plants compete for sunlight 246

Leaf angle and leaf movement can control light absorption 247

Plants acclimate and adapt to sun and shade environments 248

Photosynthetic Responses to Light by the Intact Leaf 249

Light-response curves reveal photosynthetic properties 249

Leaves must dissipate excess light energy 251

Absorption of too much light can lead to photoinhibition 253

Photosynthetic Responses to Temperature 254

Leaves must dissipate vast quantities of heat 254

Photosynthesis is temperature sensitive 255

There is an optimal temperature for photosynthesis 256

Photosynthetic Responses to Carbon Dioxide 256

Atmospheric CO₂ concentration keeps rising 257

CO₂ diffusion to the chloroplast is essential to photosynthesis 258

Patterns of light absorption generate gradients of CO₂ fixation 259

CO₂ imposes limitations on photosynthesis 260

How will photosynthesis and respiration change in the future under elevated CO₂ conditions? 261

Identifying Different Photosynthetic Pathways 263

How do we measure the stable carbon isotopes of plants? 263

Why are there carbon isotope ratio variations in plants? 264

SUMMARY 266

CHAPTER 10 Translocation in the Phloem 271**Pathways of Translocation 272**

- Sugar is translocated in phloem sieve elements 273
- Mature sieve elements are living cells specialized for translocation 273
- Large pores in cell walls are the prominent feature of sieve elements 274
- Damaged sieve elements are sealed off 274
- Companion cells aid the highly specialized sieve elements 276

Patterns of Translocation: Source to Sink 276**Materials Translocated in the Phloem 277**

- Phloem sap can be collected and analyzed 278
- Sugars are translocated in nonreducing form 279
- Other solutes are translocated in the phloem 280

Rates of Movement 280**The Pressure-Flow Model, a Passive Mechanism for Phloem Transport 281**

- An osmotically-generated pressure gradient drives translocation in the pressure-flow model 281
- The predictions of mass flow have been confirmed 282
- Sieve plate pores are open channels 283
- There is no bidirectional transport in single sieve elements 284
- The energy requirement for transport through the phloem pathway is small 284
- Positive pressure gradients exist in the phloem sieve elements 284
- Does translocation in gymnosperms involve a different mechanism? 285

Phloem Loading 285

- Phloem loading can occur via the apoplast or symplast 285
- Abundant data support the existence of apoplastic loading in some species 286
- Sucrose uptake in the apoplastic pathway requires metabolic energy 286

Phloem loading in the apoplastic pathway involves a sucrose–H⁺ symporter 287

Phloem loading is symplastic in some species 288

The polymer-trapping model explains symplastic loading in plants with intermediary cells 288

Phloem loading is passive in a number of tree species 289

The type of phloem loading is correlated with a number of significant characteristics 290

Phloem Unloading and Sink-to-Source Transition 291

Phloem unloading and short-distance transport can occur via symplastic or apoplastic pathways 291

Transport into sink tissues requires metabolic energy 292

The transition of a leaf from sink to source is gradual 292

Photosynthate Distribution: Allocation and Partitioning 294

Allocation includes storage, utilization, and transport 294

Various sinks partition transport sugars 295

Source leaves regulate allocation 295

Sink tissues compete for available translocated photosynthate 296

Sink strength depends on sink size and activity 296

The source adjusts over the long term to changes in the source-to-sink ratio 297

The Transport of Signaling Molecules 297

Turgor pressure and chemical signals coordinate source and sink activities 297

Proteins and RNAs function as signal molecules in the phloem to regulate growth and development 298

SUMMARY 299

CHAPTER 11 Respiration and Lipid Metabolism 305

Overview of Plant Respiration 305

Glycolysis 309

- Glycolysis metabolizes carbohydrates from several sources 309
- The energy-conserving phase of glycolysis extracts usable energy 310
- Plants have alternative glycolytic reactions 310
- In the absence of oxygen, fermentation regenerates the NAD^+ needed for glycolysis 311
- Plant glycolysis is controlled by its products 312

The Oxidative Pentose Phosphate Pathway 312

- The oxidative pentose phosphate pathway produces NADPH and biosynthetic intermediates 314
- The oxidative pentose phosphate pathway is redox-regulated 314

The Citric Acid Cycle 315

- Mitochondria are semiautonomous organelles 315
- Pyruvate enters the mitochondrion and is oxidized via the citric acid cycle 316
- The citric acid cycle of plants has unique features 317

Mitochondrial Electron Transport and ATP Synthesis 317

- The electron transport chain catalyzes a flow of electrons from NADH to O_2 318
- The electron transport chain has supplementary branches 320
- ATP synthesis in the mitochondrion is coupled to electron transport 320
- Transporters exchange substrates and products 322

- Aerobic respiration yields about 60 molecules of ATP per molecule of sucrose 322
- Several subunits of respiratory complexes are encoded by the mitochondrial genome 324
- Plants have several mechanisms that lower the ATP yield 324
- Short-term control of mitochondrial respiration occurs at different levels 326
- Respiration is tightly coupled to other pathways 327

Respiration in Intact Plants and Tissues 327

- Plants respire roughly half of the daily photosynthetic yield 328
- Respiration operates during photosynthesis 329
- Different tissues and organs respire at different rates 329
- Environmental factors alter respiration rates 329

Lipid Metabolism 330

- Fats and oils store large amounts of energy 331
- Triacylglycerols are stored in oil bodies 331
- Polar glycerolipids are the main structural lipids in membranes 332
- Fatty acid biosynthesis consists of cycles of two-carbon addition 334
- Glycerolipids are synthesized in the plastids and the ER 335
- Lipid composition influences membrane function 336
- Membrane lipids are precursors of important signaling compounds 336
- Storage lipids are converted into carbohydrates in germinating seeds 336

SUMMARY 338

CHAPTER 12 Assimilation of Mineral Nutrients 343**Nitrogen in the Environment 344**

Nitrogen passes through several forms in a biogeochemical cycle 344

Unassimilated ammonium or nitrate may be dangerous 346

Nitrate Assimilation 346

Many factors regulate nitrate reductase 347

Nitrite reductase converts nitrite to ammonium 347

Both roots and shoots assimilate nitrate 348

Ammonium Assimilation 348

Converting ammonium to amino acids requires two enzymes 348

Ammonium can be assimilated via an alternative pathway 350

Transamination reactions transfer nitrogen 350

Asparagine and glutamine link carbon and nitrogen metabolism 350

Amino Acid Biosynthesis 351**Biological Nitrogen Fixation 351**

Free-living and symbiotic bacteria fix nitrogen 351

Nitrogen fixation requires anaerobic conditions 352

Symbiotic nitrogen fixation occurs in specialized structures 354

Establishing symbiosis requires an exchange of signals 354

Nod factors produced by bacteria act as signals for symbiosis 354

Nodule formation involves phytohormones 355

The nitrogenase enzyme complex fixes N_2 357

Amides and ureides are the transported forms of nitrogen 358

Sulfur Assimilation 358

Sulfate is the absorbed form of sulfur in plants 358

Sulfate assimilation requires the reduction of sulfate to cysteine 359

Sulfate assimilation occurs mostly in leaves 360

Methionine is synthesized from cysteine 360

Phosphate Assimilation 360**Cation Assimilation 361**

Cations form noncovalent bonds with carbon compounds 361

Roots modify the rhizosphere to acquire iron 362

Iron forms complexes with carbon and phosphate 363

Oxygen Assimilation 363**The Energetics of Nutrient Assimilation 364****SUMMARY 365****CHAPTER 13 Secondary Metabolites and Plant Defense 369****Secondary Metabolites 370**

Secondary metabolites defend plants against herbivores and pathogens 370

Secondary metabolites are divided into three major groups 370

Terpenes 370

Terpenes are formed by the fusion of five-carbon isoprene units 370

There are two pathways for terpene biosynthesis 370

IPP and its isomer combine to form larger terpenes 371

Some terpenes have roles in growth and development 373

Terpenes defend many plants against herbivores 373

Phenolic Compounds 374

Phenylalanine is an intermediate in the biosynthesis of most plant phenolics 375

Ultraviolet light activates some simple phenolics 377

The release of phenolics into the soil may limit the growth of other plants 377

Lignin is a highly complex phenolic macromolecule 377

There are four major groups of flavonoids 378
 Anthocyanins are colored flavonoids that attract animals 378
 Flavones and flavonols may protect against damage by ultraviolet light 379
 Isoflavonoids have widespread pharmacological activity 379
 Tannins deter feeding by herbivores 380

Nitrogen-Containing Compounds 381

Alkaloids have dramatic physiological effects on animals 381
 Cyanogenic glycosides release the poison hydrogen cyanide 384
 Glucosinolates release volatile toxins 385
 Nonprotein amino acids are toxic to herbivores 385

Induced Plant Defenses against Insect Herbivores 386

Plants can recognize specific components of insect saliva 386
 Jasmonic acid activates many defensive responses 387
 Some plant proteins inhibit herbivore digestion 389

Damage by insect herbivores induces systemic defenses 389
 Herbivore-induced volatiles have complex ecological functions 389
 Insects have developed strategies to cope with plant defenses 391

Plant Defenses against Pathogens 391

Pathogens have developed various strategies to invade host plants 391
 Some antimicrobial compounds are synthesized before pathogen attack 392
 Infection induces additional antipathogen defenses 392
 Phytoalexins often increase after pathogen attack 393
 Some plants recognize specific pathogen-derived substances 393
 Exposure to elicitors induces a signal transduction cascade 394
 A single encounter with a pathogen may increase resistance to future attacks 394
 Interactions of plants with nonpathogenic bacteria can trigger induced systemic resistance 395

SUMMARY 396

UNIT III ■ Growth and Development 401

CHAPTER 14 ■ Signal Transduction 403

Signal Transduction in Plant and Animal Cells 404

Plants and animals have similar transduction components 404
 Receptor kinases can initiate a signal transduction cascade 406
 Plants signal transduction components have evolved from both prokaryotic and eukaryotic ancestors 406
 Signals are perceived at many locations within plant cells 408
 Plant signal transduction often involves inactivation of repressor proteins 409
 Protein degradation is a common feature in plant signaling pathways 411

Several plant hormone receptors encode components of the ubiquitination machinery 413
 Inactivation of repressor proteins results in a gene expression response 414
 Plants have evolved mechanisms for switching off or attenuating signaling responses 414
 Cross-regulation allows signal transduction pathways to be integrated 416

Signal Transduction in Space and Time 418

Plant signal transduction occurs over a wide range of distances 418
 The timescale of plant signal transduction ranges from seconds to years 419

SUMMARY 421

CHAPTER 15 Cell Walls: Structure, Biogenesis, and Expansion 425

The Structure and Synthesis of Plant Cell Walls 426

- Plant cell walls have varied architecture 426
- The primary cell wall is composed of cellulose microfibrils embedded in a polysaccharide matrix 428
- Cellulose microfibrils are synthesized at the plasma membrane 430
- Matrix polymers are synthesized in the Golgi apparatus and secreted via vesicles 433
- Hemicelluloses are matrix polysaccharides that bind to cellulose 433
- Pectins are hydrophilic gel-forming components of the matrix 434
- Structural proteins become cross-linked in the wall 437
- New primary walls are assembled during cytokinesis 437

Secondary walls form in some cells after expansion ceases 438

Patterns of Cell Expansion 441

- Microfibril orientation influences growth directionality of cells with diffuse growth 441
- Cortical microtubules influence the orientation of newly deposited microfibrils 443

The Rate of Cell Elongation 443

- Stress relaxation of the cell wall drives water uptake and cell elongation 445
- Acid-induced growth and wall stress relaxation are mediated by expansins 446
- Many structural changes accompany the cessation of wall expansion 448

SUMMARY 448

CHAPTER 16 Growth and Development 453

Overview of Plant Growth and Development 454

- Sporophytic development can be divided into three major stages 455

Embryogenesis: The Origins of Polarity 456

- Embryogenesis differs between dicots and monocots, but also features common fundamental processes 456
- Apical–basal polarity is established early in embryogenesis 457
- Position-dependent signaling guides embryogenesis 458
- Auxin may function as a mobile chemical signal during embryogenesis 460
- Mutant analysis has helped identify genes essential for embryo organization 461
- The GNOM protein establishes a polar distribution of auxin efflux proteins 463
- MONOPTEROS encodes a transcription factor that is activated by auxin 463
- Radial patterning guides formation of tissue layers 464

The differentiation of cortical and endodermal cells involves the intercellular movement of a transcription factor 465

Many developmental processes involve the intercellular movement of macromolecules 467

Meristematic Tissues: Foundations for Indeterminate Growth 468

The root and shoot apical meristems use similar strategies to enable indeterminate growth 469

The Root Apical Meristem 469

- The root tip has four developmental zones 469
- The origin of different root tissues can be traced to specific initial cells 470
- Cell ablation experiments implicate directional signaling processes in determination of cell identity 471
- Auxin contributes to the formation and maintenance of the RAM 471
- Responses to auxin depend on specific transcription factors 472
- Cytokinin activity in the RAM is required for root development 473

The Shoot Apical Meristem 474

- The shoot apical meristem has distinct zones and layers 474
- Shoot tissues are derived from several discrete sets of apical initials 475
- The locations of PIN proteins influence SAM formation 476
- Embryonic SAM formation requires the coordinated expression of transcription factors 477
- Negative feedback limits apical meristem size 478
- Similar mechanisms maintain initials in the RAM and in the SAM 479

Vegetative Organogenesis 480

- Localized zones of auxin accumulation promote leaf initiation 480
- Spatially regulated gene expression determines the planar form of the leaf 481
- Distinct mechanisms initiate roots and shoots 483

Senescence and Programmed Cell Death 484

- Leaf senescence is adaptive and strictly regulated 484
- Plants exhibit various types of senescence 485
- Senescence involves the ordered degradation of potentially phototoxic chlorophyll 487
- Programmed cell death is a specialized type of senescence 487

SUMMARY 488

CHAPTER 17 Phytochrome and Light Control of Plant Development 493

The Photochemical and Biochemical Properties of Phytochrome 494

- Phytochrome can interconvert between Pr and Pfr forms 496
- Pfr is the physiologically active form of phytochrome 496

Characteristics of Phytochrome-Induced Responses 497

- Phytochrome responses vary in lag time and escape time 497
- Phytochrome responses can be distinguished by the amount of light required 497
- Very low-fluence responses are nonphotoreversible 497
- Low-fluence responses are photoreversible 498
- High-irradiance responses are proportional to the irradiance and the duration 499

Structure and Function of Phytochrome Proteins 499

- Phytochrome has several important functional domains 500
- Phytochrome is a light-regulated protein kinase 501
- Pfr is partitioned between the cytosol and the nucleus 501
- Phytochromes are encoded by a multigene family 502

Genetic Analysis of Phytochrome Function 503

- Phytochrome A mediates responses to continuous far-red light 504
- Phytochrome B mediates responses to continuous red or white light 504
- Roles for phytochromes C, D, and E are emerging 504
- Phy gene family interactions are complex 504
- PHY gene functions have diversified during evolution 505

Phytochrome Signaling Pathways 505

- Phytochrome regulates membrane potentials and ion fluxes 506
- Phytochrome regulates gene expression 506
- Phytochrome interacting factors (PIFs) act early in phy signaling 507
- Phytochrome associates with protein kinases and phosphatases 507
- Phytochrome-induced gene expression involves protein degradation 508

Circadian Rhythms 509

- The circadian oscillator involves a transcriptional negative feedback loop 510

Ecological Functions 512

- Phytochrome enables plant adaptation to changes in light quality 512

XXVIII TABLE OF CONTENTS

Decreasing the R:FR ratio causes elongation in sun plants 512
Small seeds typically require a high R:FR ratio for germination 513
Reducing shade avoidance responses can improve crop yields 514

Phytochrome responses show ecotypic variation 515
Phytochrome action can be modulated 515

SUMMARY 516

CHAPTER 18 Blue-Light Responses: Morphogenesis and Stomatal Movements 521

The Photophysiology of Blue-Light Responses 522

Blue light stimulates asymmetric growth and bending 523
Blue light rapidly inhibits stem elongation 523
Blue light stimulates stomatal opening 524
Blue light activates a proton pump at the guard cell plasma membrane 527
Blue-light responses have characteristic kinetics and lag times 528
Blue light regulates the osmotic balance of guard cells 528
Sucrose is an osmotically active solute in guard cells 530

The Regulation of Blue Light–Stimulated Responses 531

Blue-Light Photoreceptors 532

Cryptochromes regulate plant development 532
Phototropins mediate blue light–dependent phototropism and chloroplast movements 533
Zeaxanthin mediates blue-light photoreception in guard cells 534
Green light reverses blue light–stimulated opening 536

SUMMARY 539

CHAPTER 19 Auxin: The First Discovered Plant Growth Hormone 545

The Emergence of the Auxin Concept 546

The Principal Auxin: Indole-3-Acetic Acid 546

IAA is synthesized in meristems and young dividing tissues 549
Multiple pathways exist for the biosynthesis of IAA 549
Seeds and storage organs contain covalently bound auxin 550
IAA is degraded by multiple pathways 550

Auxin Transport 551

Polar transport requires energy and is gravity independent 552
Chemiosmotic potential drives polar transport 553
PIN and ABCB transporters regulate cellular auxin homeostasis 555
Auxin influx and efflux can be chemically inhibited 556
Auxin transport is regulated by multiple mechanisms 558

Auxin Signal Transduction Pathways 560

The principal auxin receptors are soluble protein heterodimers 561
Auxin-induced genes are negatively regulated by AUX/IAA proteins 561
Auxin binding to a TIR1/AFB-AUX/IAA heterodimer stimulates AUX/IAA destruction 562
Auxin-induced genes fall into two classes: early and late 562
Rapid, nontranscriptional auxin responses appear to involve a different receptor protein 562

Actions of Auxin: Cell Elongation 562

Auxins promote growth in stems and coleoptiles, while inhibiting growth in roots 563
The outer tissues of dicot stems are the targets of auxin action 563
The minimum lag time for auxin-induced elongation is ten minutes 565

Auxin rapidly increases the extensibility of the cell wall 565

Auxin-induced proton extrusion increases cell extension 565

Auxin-induced proton extrusion involves activation and protein mobilization 566

Actions of Auxin: Plant Tropisms 566

Phototropism is mediated by the lateral redistribution of auxin 566

Gravitropism involves lateral redistribution of auxin 568

Dense plastids serve as gravity sensors 569

Gravity sensing may involve pH and calcium ions (Ca^{2+}) as second messengers 571

Auxin is redistributed laterally in the root cap 572

Developmental Effects of Auxin 573

Auxin regulates apical dominance 574

Auxin transport regulates floral bud development and phyllotaxy 576

Auxin promotes the formation of lateral and adventitious roots 576

Auxin induces vascular differentiation 576

Auxin delays the onset of leaf abscission 577

Auxin promotes fruit development 577

Synthetic auxins have a variety of commercial uses 578

SUMMARY 578

CHAPTER 20

Gibberellins: Regulators of Plant Height and Seed Germination 583

Gibberellins: Their Discovery and Chemical Structure 584

Gibberellins were discovered by studying a disease of rice 584

Gibberellic acid was first purified from *Gibberella* culture filtrates 584

All gibberellins are based on an *ent*-gibberellane skeleton 585

Effects of Gibberellins on Growth and Development 586

Gibberellins promote seed germination 586

Gibberellins can stimulate stem and root growth 586

Gibberellins regulate the transition from juvenile to adult phases 587

Gibberellins influence floral initiation and sex determination 588

Gibberellins promote pollen development and tube growth 588

Gibberellins promote fruit set and parthenocarpy 588

Gibberellins promote early seed development 588

Commercial uses of gibberellins and GA biosynthesis inhibitors 588

Biosynthesis and Deactivation of Gibberellins 589

Gibberellins are synthesized via the terpenoid pathway 589

Some enzymes in the GA pathway are highly regulated 591

Gibberellin regulates its own metabolism 592

GA biosynthesis occurs at multiple plant organs and cellular sites 592

Environmental conditions can influence GA biosynthesis 593

GA_1 and GA_4 have intrinsic bioactivity for stem growth 594

Plant height can be genetically engineered 595

Dwarf mutants often show other phenotypic defects 595

Auxins can regulate GA biosynthesis 595

Gibberellin Signaling: Significance of Response Mutants 596

GID1 encodes a soluble GA receptor 596

DELLA-domain proteins are negative regulators of GA response 600

Mutation of negative regulators of GA may produce slender or dwarf phenotypes 600

Gibberellins signal the degradation of negative regulators of GA response 601

F-box proteins target DELLA domain proteins for degradation 601

Negative regulators with DELLA domains have agricultural importance 602

Gibberellin Responses: Early Targets of DELLA Proteins 602

DELLA proteins can activate or suppress gene expression 603

DELLA proteins regulate transcription by interacting with other proteins such as phytochrome-interacting factors 603

Gibberellin Responses: The Cereal Aleurone Layer 605

GA is synthesized in the embryo 605

Aleurone cells may have two types of GA receptors 605

Gibberellins enhance the transcription of α -amylase mRNA 605

GAMYB is a positive regulator of α -amylase transcription 607

DELLA-domain proteins are rapidly degraded 607

Gibberellin Responses: Anther Development and Male Fertility 607

GAMYB regulates male fertility 609

Events downstream of GAMYB in rice aleurone and anthers are quite different 611

MicroRNAs regulate MYBs after transcription in anthers but not in aleurone 611

Gibberellin Responses: Stem Growth 612

Gibberellins stimulate cell elongation and cell division 612

GAs regulate the transcription of cell cycle kinases 613

Reducing GA sensitivity may prevent crop losses 613

SUMMARY 614

CHAPTER 21 Cytokinins: Regulators of Cell Division 621

Cell Division and Plant Development 622

Differentiated plant cells can resume division 622

Diffusible factors control cell division 622

Plant tissues and organs can be cultured 622

The Discovery, Identification, and Properties of Cytokinins 623

Kinetin was discovered as a breakdown product of DNA 623

Zeatin was the first natural cytokinin discovered 623

Some synthetic compounds can mimic cytokinin action 624

Cytokinins occur in both free and bound forms 625

Some plant pathogenic bacteria, fungi, insects, and nematodes secrete free cytokinins 625

Biosynthesis, Metabolism, and Transport of Cytokinins 625

Crown gall cells have acquired a gene for cytokinin synthesis 626

IPT catalyzes the first step in cytokinin biosynthesis 628

Cytokinins can act both as long distance and local signals 628

Cytokinins are rapidly metabolized by plant tissues 628

Cellular and Molecular Modes of Cytokinin Action 629

A cytokinin receptor related to bacterial two-component receptors has been identified 629

Cytokinins increase expression of the type-A response regulator genes via activation of the type-B ARR genes 630

Histidine phosphotransfer proteins are also involved in cytokinin signaling 632

The Biological Roles of Cytokinins 632

Cytokinins promote shoot growth by increasing cell proliferation in the shoot apical meristem 632

Cytokinins interact with other hormones and with several key transcription factors 634

Cytokinins inhibit root growth by promoting the exit of cells from the root apical meristem 635

Cytokinins regulate specific components of the cell cycle 636

The auxin:cytokinin ratio regulates morphogenesis in cultured tissues 637

Cytokinins modify apical dominance and promote lateral bud growth 638

Cytokinins delay leaf senescence 638

Cytokinins promote movement of nutrients 639

Cytokinins affect light signaling via phytochrome 640
 Cytokinins regulate vascular development 641
 Manipulation of cytokinins to alter agriculturally important traits 641

Cytokinins are involved in the formation of nitrogen-fixing nodules in legumes 641

SUMMARY 643

CHAPTER 22 Ethylene: The Gaseous Hormone 649

Structure, Biosynthesis, and Measurement of Ethylene 650

Regulated biosynthesis determines the physiological activity of ethylene 650
 Ethylene biosynthesis is promoted by several factors 652
 Ethylene biosynthesis can be elevated through a stabilization of ACC synthase protein 652
 Various inhibitors can block ethylene biosynthesis 653

Ethylene Signal Transduction Pathways 653

Ethylene receptors are related to bacterial two-component system histidine kinases 654
 High-affinity binding of ethylene to its receptor requires a copper cofactor 655
 Unbound ethylene receptors are negative regulators of the response pathway 655
 A serine/threonine protein kinase is also involved in ethylene signaling 657
 EIN2 encodes a transmembrane protein 657

Ethylene Regulation of Gene Expression 657

Specific transcription factors are involved in ethylene-regulated gene expression 657
 Genetic epistasis reveals the order of the ethylene signaling components 658

Developmental and Physiological Effects of Ethylene 659

Ethylene promotes the ripening of some fruits 659
 Fruits that respond to ethylene exhibit a climacteric 659
 The receptors of *never-ripe* mutants of tomato fail to bind ethylene 660
 Leaf epinasty results when ACC from the root is transported to the shoot 660
 Ethylene induces lateral cell expansion 661
 There are two distinct phases to growth inhibition by ethylene 662
 The hooks of dark-grown seedlings are maintained by ethylene production 662
 Ethylene breaks seed and bud dormancy in some species 663
 Ethylene promotes the elongation growth of submerged aquatic species 663
 Ethylene induces the formation of roots and root hairs 664
 Ethylene regulates flowering and sex determination in some species 664
 Ethylene enhances the rate of leaf senescence 664
 Ethylene mediates some defense responses 665
 Ethylene acts on the abscission layer 665
 Ethylene has important commercial uses 667

SUMMARY 668

CHAPTER 23 Abscisic Acid: A Seed Maturation and Stress-Response Hormone 673

Occurrence, Chemical Structure, and Measurement of ABA 674

The chemical structure of ABA determines its physiological activity 674
 ABA is assayed by biological, physical, and chemical methods 674

Biosynthesis, Metabolism, and Transport of ABA 674

ABA is synthesized from a carotenoid intermediate 674
 ABA concentrations in tissues are highly variable 676
 ABA is translocated in vascular tissue 677

ABA Signal Transduction Pathways 678

Receptor candidates include diverse classes of proteins 678
 Secondary messengers function in ABA signaling 680
 Ca²⁺-dependent and Ca²⁺-independent pathways mediate ABA signaling 680
 ABA-induced lipid metabolism generates second messengers 681
 Protein kinases and phosphatases regulate important steps in ABA signaling 682
 PP2Cs interact directly with the PYR/PYL/RCAR family of ABA receptors 683
 ABA shares signaling intermediates with other hormonal pathways 683

ABA Regulation of Gene Expression 683

Gene activation by ABA is mediated by transcription factors 684

Developmental and Physiological Effects of ABA 684

ABA regulates seed maturation 684
 ABA inhibits precocious germination and vivipary 685
 ABA promotes seed storage reserve accumulation and desiccation tolerance 686
 Seed dormancy can be regulated by ABA and environmental factors 686
 Seed dormancy is controlled by the ratio of ABA to GA 687
 ABA inhibits GA-induced enzyme production 688
 ABA promotes root growth and inhibits shoot growth at low water potentials 688
 ABA promotes leaf senescence independently of ethylene 689
 ABA accumulates in dormant buds 689
 ABA closes stomata in response to water stress 690
 ABA regulates ion channels and the plasma membrane ATPase in guard cells 690

SUMMARY 693

CHAPTER 24 **Brassinosteroids: Regulators of Cell Expansion and Development 699**

Brassinosteroid Structure, Occurrence, and Genetic Analysis 700

BR-deficient mutants are impaired in photomorphogenesis 701

The Brassinosteroid Signaling Pathway 703

BR-insensitive mutants identified the BR cell surface receptor 703
 Phosphorylation activates the BRI1 receptor 704
 BIN2 is a repressor of BR-induced gene expression 704
 BES1/BZR1 regulate gene expression 706

Biosynthesis, Metabolism, and Transport of Brassinosteroids 706

Brassinolide is synthesized from campesterol 707
 Catabolism and negative feedback contribute to BR homeostasis 708

Brassinosteroids act locally near their sites of synthesis 710

Brassinosteroids: Effects on Growth and Development 710

BRs promote both cell expansion and cell division in shoots 711
 BRs both promote and inhibit root growth 712
 BRs promote xylem differentiation during vascular development 713
 BRs are required for the growth of pollen tubes 714
 BRs promote seed germination 714

Prospective Uses of Brassinosteroids in Agriculture 714

SUMMARY 715

CHAPTER 25 The Control of Flowering 719

Floral Meristems and Floral Organ Development 720

The shoot apical meristem in *Arabidopsis* changes with development 721

The four different types of floral organs are initiated as separate whorls 721

Two major types of genes regulate floral development 722

Meristem identity genes regulate meristem function 722

Homeotic mutations led to the identification of floral organ identity genes 723

Three types of homeotic genes control floral organ identity 723

The ABC model explains the determination of floral organ identity 724

Floral Evocation: Integrating Environmental Cues 725

The Shoot Apex and Phase Changes 726

Plant development has three phases 726

Juvenile tissues are produced first and are located at the base of the shoot 727

Phase changes can be influenced by nutrients, gibberellins, and other signals 728

Competence and determination are two stages in floral evocation 728

Circadian Rhythms: The Clock Within 730

Circadian rhythms exhibit characteristic features 730

Phase shifting adjusts circadian rhythms to different day–night cycles 732

Phytochromes and cryptochromes entrain the clock 732

Photoperiodism: Monitoring Day Length 732

Plants can be classified according to their photoperiodic responses 732

The leaf is the site of perception of the photoperiodic signal 734

Plants monitor day length by measuring the length of the night 734

Night breaks can cancel the effect of the dark period 735

The circadian clock and photoperiodic timekeeping 736

The coincidence model is based on oscillating light sensitivity 737

The coincidence of *CONSTANS* expression and light promotes flowering in LDPs 737

SDPs use a coincidence mechanism to inhibit flowering in long days 739

Phytochrome is the primary photoreceptor in photoperiodism 739

A blue-light photoreceptor regulates flowering in some LDPs 740

Vernalization: Promoting Flowering with Cold 741

Vernalization results in competence to flower at the shoot apical meristem 742

Vernalization can involve epigenetic changes in gene expression 742

A range of vernalization pathways may have evolved 743

Long-Distance Signaling Involved in Flowering 744

The floral stimulus is transported in the phloem 744

Grafting studies have provided evidence for a transmissible floral stimulus 744

The Discovery of Florigen 745

The *Arabidopsis* protein FLOWERING LOCUS T is florigen 746

Gibberellins and ethylene can induce flowering 747

Climate change has already caused measurable changes in flowering time of wild plants 748

The transition to flowering involves multiple factors and pathways 748

SUMMARY 749

CHAPTER 26 Responses and Adaptations to Abiotic Stress 755**Adaptation and Phenotypic Plasticity 756**

Adaptations involve genetic modification 756
 Phenotypic plasticity allows plants to respond to environmental fluctuations 756

The Abiotic Environment and its Biological Impact on Plants 756

Climate and soil influence plant fitness 757
 Imbalances in abiotic factors have primary and secondary effects on plants 757

Water Deficit and Flooding 757

Soil water content and the relative humidity of the atmosphere determine the water status of the plant 758
 Water deficits cause cell dehydration and an inhibition of cell expansion 759
 Flooding, soil compaction, and O₂ deficiency are related stresses 759

Imbalances in Soil Minerals 760

Soil mineral content can result in plant stress in various ways 760
 Soil salinity occurs naturally and as the result of improper water management practices 761
 The toxicity of high Na⁺ and Cl⁻ in the cytosol is due to their specific ion effects 761

Temperature Stress 762

High temperatures are most damaging to growing, hydrated tissues 762
 Temperature stress can result in damaged membranes and enzymes 762
 Temperature stress can inhibit photosynthesis 763
 Low temperatures above freezing can cause chilling injury 764
 Freezing temperatures cause ice crystal formation and dehydration 764

High Light Stress 764

Photoinhibition by high light leads to the production of destructive forms of oxygen 764

Developmental and Physiological Mechanisms that Protect Plants against Environmental Extremes 765

Plants can modify their life cycles to avoid abiotic stress 765
 Phenotypic changes in leaf structure and behavior are important stress responses 765
 The ratio of root-to-shoot growth increases in response to water deficit 769
 Plants can regulate stomatal aperture in response to dehydration stress 769
 Plants adjust osmotically to drying soil by accumulating solutes 769
 Submerged organs develop aerenchyma tissue in response to hypoxia 770
 Plants have evolved two different strategies to protect themselves from toxic ions: exclusion and internal tolerance 772
 Chelation and active transport contribute to internal tolerance 773
 Many plants have the capacity to acclimate to cold temperatures 773
 Plants survive freezing temperatures by limiting ice formation 774
 The lipid composition of membranes affects their response to temperature 775
 Plant cells have mechanisms that maintain protein structure during temperature stress 776
 Scavenging mechanisms detoxify reactive oxygen species 776
 Metabolic shifts enable plants to cope with a variety of abiotic stresses 777

SUMMARY 778**APPENDIX ONE A1-1****APPENDIX TWO A2-1****APPENDIX THREE A3-1****GLOSSARY G-1****AUTHOR INDEX AI-1****SUBJECT INDEX SI-1**